

RESEARCH PAPER

A Preliminary Checklist of Auchenorrhyncha (Hemiptera) from Bangladesh

Mohammad Atikur Rahman^{1*}, Pijush Kanti Jhan¹, Md. Mohasin Hussain Khan¹, S. M. Hemayet Jahan¹, Rahagir Alam¹

¹ Department of Entomology, Patuakhali Science and Technology University, Dumki, Patuakhali-8602, Bangladesh

ARTICLE HISTORY

Received: October 10, 2016
Revised : November 8, 2016
Accepted: November 22, 2016
Published: December 31, 2016

*Corresponding author
E-mail:atikentom@pstu.ac.bd

ABSTRACT

A preliminary checklist of Auchenorrhyncha (Hemiptera) from Bangladesh is provided, based primarily on literature records from 1758 to the present. In total 11 families, 34 tribes, 55 genera and 74 species are recorded with taxonomic details. Among 25 genera of Cicadomorpha, 7 genera are firstly recognized herein: *Goniagnathus* Fieber, 1866; *Hecalus* Walker, 1858 ; *Exitianus* Ball, 1929; *Kolla* Distant, 1908; *Rhytidodus* Fieber, 1868; *Scaphoidella* Vilbaste, 1968; *Penthimia* Germar, 1821 and 8 species of them are recorded for the first time in Bangladesh: *Goniagnathus punctifer* (Walker, 1858); *Hecalus porrectus* (Walker, 1858); *Scaphoidella* sp. Vilbaste, 1968; *Penthimia picta* Provancher, 1872; *Penthimia nigra* Goeze, 1778; *Kolla bataviae* (Bierman, 1910); *Rhytidodus decimusquartus* Schrank, 1776; *Exitianus indicus* Distant, 1908. In the Fulgoromorpha, family Derbidae Spinola, 1839, and 3 genera are also firstly recognized herein: *Tagosodes* Asche & Wilson, 1990; *Tropidocephala* Stål, 1853; *Raivuna* Fennah, 1978 and 4 species of them are recorded for the first time in Bangladesh: *Deferunda qiana* (Chen and He, 2010); *Tagosodes cubanus* (Crawford, 1914); *Tropidocephala festiva* Distant, 1906; *Raivuna sinica* Walter, 1851.

Key words: Fulgoromorpha, Cicadomorpha, planthopper, leafhopper, checklist, Bangladesh

Introduction

The species of Auchenorrhyncha are a great concern to agriculture and forestry because several species are significant pests of a number of important crop plants, and they reach high population densities followed by reducing crop production, damaging crops and/or they act as a vector of some bacteria, phytoplasma, or viral pathogens. Bangladesh is a humid and subtropical country favoring luxuriant growth of various insect species with rich diversity. In the beginning of 20th century, before the birth of Bangladesh, Distant (1900-1920) worked with the insect fauna of British India where he collected very few species of Auchenorrhyncha from the southeastern border of present Bangladesh. After that, a very few taxonomic work had been done in Bangladesh. Therefore, collection, identification and records of such insect species are barely necessary in Bangladesh. Some works on Auchenorrhyncha species in Bangladesh have been done by the following authors: Ahmed *et al.*

(1970), Lallemand (1963), Hope (1843), Constant (2010), Stål (1863), Ishihara and Lippole (1969), Chen & He (2010), and Distant (1906). A comparative detailed study of the Oriental fulgoroida was undertaken by Distant (1906-1916) who erected twelve families in his work on Rhynchota in the series of publication on the Fauna of British India including Ceylon and Burma where he recorded some fulgoroids from southeast and northeast part of Bangladesh. Recently Jhan *et al.* (2016) has worked on leafhopper and planthopper from Southern and South eastern part of Bangladesh, and reported few new records.

In the current study a list of Auchenorrhyncha species recorded from Bangladesh has been compiled, which is based on literature records and the addition of some newly recorded species. Much basic taxonomic work is needed in Bangladesh and it is hoped that this preliminary checklist will facilitate further studies on Auchenorrhyncha of Bangladesh.

Checklist

Infraorder Cicadomorpha

Family Cicadidae

Tribe Platycleurini

***Pycna repanda repanda* (Linnaeus, 1758)**

Synonym:

Cicada repanda Linnaeus, 1758: 436

Tettigonia repanda (Linnaeus, 1758)

Platypleura phalaenoides Walker, 1850

Fidicina repanda (Linnaeus, 1758)

Platypleura interna Walker, 1852

Oxypleura repanda Dohrn, 1859

Platypleura congrex Butler, 1874

Platypleura repanda (Linnaeus, 1758)

Pycnarepanda (Linnaeus, 1758)

Platypleura nepanda (Linnaeus, 1758) (sic)

Distribution: India, Bangladesh, Japan, Europe, Taiwan, China, Pakistan, Indonesia, Nepal, Myanmar, Bhutan, Sri Lanka.

Tribe Tosenini

***Tosena dives* (Westwood, 1842)**

Synonym:

Cicada dives Westwood, 1842: 98 (25)

Huechys transversa Walker, 1858

Tosena dives (Westwood, 1842)

Distribution: India, Bangladesh, Nepal.

***Tosena mearesiana* (Westwood, 1842)**

Synonym:

Cicada mearesiana Westwood, 1842: 98

Distribution: India, Bangladesh, Nepal.

***Tosena melanopteryx* (Kirkaldy, 1909)**

Synonym:

Cicada melanoptera White, 1846:331

Tosena albata melanopteryx Kirkaldy, 1909: 391

Tosena fasciata Moulton, 1923

Tosena melanoptera White 1846

Tosena melanopteryx (Kirkaldy, 1909) (sic)

Tosena melanopteryx (Kirkaldy, 1909)

Distribution: India, Bangladesh, Nepal, Myanmar, Thailand, Vietnam, China, Laos.

Tribe Cryptotympanini

***Cryptotympana corvus* (Walker, 1850)**

Synonym:

Fidicina corvus Walker, 1850: 86

Cryptotympana corva Stål, 1862

Cryptotympana invariants Atkinson, 1886

Fidicina invariants Walker, 1858

Distribution: India, Bangladesh, Nepal, Bhutan.

***Cryptotympana recta* (Walker, 1850)**

Synonym:

Fidicina recta Walker, 1850: 79

Distribution: India, Bangladesh, China, Malaysia, Laos, Thailand, Vietnam, Cambodia, Sumatra.

Tribe Dundubiini

***Dundubia ensifera* Bloem & Duffels, 1976**

Dundubia ensifera Bloem & Duffels, 1976: 146.

Distribution: India, Bangladesh, Bhutan, Nepal, Sri Lanka.

***Haphsa bindusara* (Distant 1881)**

Synonym:

Pomponia bindusara Distant, 1881:642

Chremistica ochracea Boulard, 2005 (sic)

Aola bindusara (Distant, 1881)

Haphsa bindusara (Distant, 1881) (sic)

Distribution: India, Bangladesh, Pakistan, Myanmar, Vietnam, China, Laos, Thailand.

***Songga scitula* (Distant, 1888)**

Synonym:

Pomponia scitula Distant, 1888: 456

Meimuna scitula (Distant, 1888)

Aola scitula (Distant, 1888)

Haphsa scitula (Distant, 1888)

Songga scitula (Distant, 1888): Lee, 2016 (p.597)

Distribution: India, Bangladesh, Myanmar, Cambodia, China, Pakistan, Thailand.

***Macrosemia saturata* (Walker, 1858)**

Synonym:

Dundubia obtecta Walker, 1850 (misid.)

Dundubia saturata Walker, 1858: 6

Cosmopsaltria saturate (Walker, 1858)

Platylomia saturata (Walker, 1858)

Meimuna saturata (Walker, 1858)

Platylomia satura (Walker, 1858) (sic)

Cosmopsaltria saturate (Walker, 1858)(sic)

Distribution: Bangladesh, India, Nepal, Bhutan, Pakistan, Malaysia, Vietnam, Indonesia.

***Platylomia vibrans* (Walker, 1850)**

Synonym:

Dundubia lateralis Walker, 1850: 61

Dundubia vibrans Walker, 1850: 54

Cosmopsaltria vibrans (Walker, 1850)

Platylomia vibrans (Walker, 1850)

Distribution: Bangladesh, India, Malaysia.

Tribe Gaeanini

***Gaeana maculata* (Drury, 1773)**

Synonym:

Cicada maculata Drury, 1773: 68

Distribution: India, Bangladesh, China, Syria, Hong Kong, Australia, Japan, Myanmar, Indonesia, Malaysia, Vietnam, Sri Lanka.

***Sulphogaeana sulphurea* (Westwood, 1839)**

Synonym:

Cicada sulphurea Westwood, 1839: 54

Cicada pulchella Westwood, 1839

Cicada pulcilla (Westwood, 1839)

Gaeana pulchella(Westwood, 1839)

Gaeana sulphurea (Westwood, 1839)

Sulphogaeana sulphura (Westwood, 1839)(sic)

Distribution: Bangladesh, India, Nepal, Bhutan, China.

Tribe Moganniini

***Mogannia funebris* Stål, 1865**

Mogannia funebris Stål, 1865: 155

Distribution: Bangladesh, India, Myanmar, Vietnam.

Tribe Cicadini

***Pomponia cinctimanus* (Walker, 1850)**

Synonym:

Dundubia cinctimanus Walker, 1850: 49

Distribution: Bangladesh, India, Vietnam

***Pomponia linearis* (Walker, 1850)**

Synonym:

Dundubia linearis Walker, 1850:48

Pomponia fusca Kato, 1956 (misapplied name)

Distribution: Bangladesh, India, Vietnam, China, Myanmar, Thailand, Laos, Nepal, Philippines, Malaysia, Singapore, Indonesia.

***Pomponia ramifera* (Walker, 1850)**

Synonym:

Dundubia ramifera Walker, 1850:53

Dundubia ramifera (Walker, 1850) (sic)

Dundubia linearis ramifera (Walker, 1850)

Pomponia fusca Kato, 1956 (misapplied name)

Distribution: Bangladesh, India, Vietnam.

***Meimuna silhetana* (Distant, 1888)**

Synonym:

Cosmopsaltria silhetana Distant, 1888: 295

Distribution: Bangladesh, India, China, Hong Kong.

Family: Cicadelidae

Tribe Cicadellini

***Cofana spectra* (Distant, 1908)**

Tettigoniella spectra Distant, 1908: 211

Distribution: Africa, Australia, Bangladesh and Oriental region.

***Kolla bataviae* (Bierman, 1910)**

Tettigonia bataviae Bierman, 1910: 51

Distribution: Bangladesh (new record).

Tribe Coelidiini

***Olidiana brevis* (Walker, 1851)**

Synonym:

Tettigonia brevis Walker, 1851:774

Olidiana brevis (Walker, 1851): McKamey, 2006: 503

Distribution: Bangladesh, Burma, China, Laos and Vietnam, Africa.

***Exitianus indicus* Distant, 1908**

Athysanus indicus Distant, 1908: 344

Distribution: India, Pakistan, Bangladesh (new record).

Tribe Chiasmini

***Nephotettix cincticeps* (Uhler, 1896)**

Selenocephalus cincticeps Uhler, 1896: 292

Distribution: Bangladesh, India, China, Japan, Korea and Taiwan. Russia.

***Nephotettix nigropictus* (Stål, 1870)**

Thamnotettix nigropictus Stål, 1870: 740

Distribution: Australia, Bangladesh, Burma, China, Guinea, India, Indonesia, Malaysia, Nepal, New Hong Kong, Pakistan, Philippines, South Vietnam, Sri Lanka and Thailand

***Nephotettix virescens* (Distant, 1908)**

Selenocephalus virescens Distant, 1908: 291

Distribution: Bangladesh, Burma, Hong Kong, India, Indonesia, Laos, Malaysia, Pakistan, Philippines, Taiwan, South Vietnam, Thailand, Sri Lanka, China and Japan.

Tribe Deltocephalini

***Maiestas dorsalis* (Motschulsky, 1859)**

Synonym:

Deltocephalus dorsalis Motschulsky, 1859: 114

Thamnotettix dorsalis Matsumura, 1907

Recilia (Inazuma) dorsalis Ishihara, 1953

Distribution: Australia, Bangladesh, Bhutan, Burma, China, India, Indonesia, Japan, Kampuchea, Korea, Laos, Malaysia, Pakistan, Philippines, Sri Lanka, Taiwan, Thailand, Vietnam, Japan, Borneo, Korea.

***Phlogotettix indicus* Rao, 1989**

Phlogotettix indicus Rao, 1989: 59-83

Distribution: India, Pakistan, Bangladesh.

Tribe Goniagnathini

***Goniagnathus punctifer* (Walker, 1858)**

Bythoscopus punctifer Walker, 1858: 104

Distribution: Bangladesh (new record).

Tribe Hecalini

***Hecalus porrectus* (Walker, 1858)**

Hecalus porrectus (Walker, 1858): 362

Distribution: Bangladesh (new record), Pakistan.

Tribe Opsiini

***Orosius orientalis* (Matsumura, 1914)**

Eutettix orientalis Matsumura, 1914: 192.

Distribution: Australia, Bangladesh, India, Japan, Myanmar and Taiwan, Fiji, New Britain, and Java.

Tribe Penthimini

***Penthimia nigra* Goeze, 1778**

Cicada nigra Goeze, 1778: 161

Distribution: Bangladesh (new record), Germany, Japan, Russia, Albania, Austria, Azerbaijan, Belgium, Bohemia, Bulgaria, Czech Republic, Czechoslovakia, Russia, France, Germany, Great Britain, Greece, Hungary, Italy, Japan, Latvia, Macedonia, Moldova, Moravia, Netherlands, Poland, Portugal, Romania, Serbia, Slovakia, Spain, Switzerland, Syria, Tunisia, Turkey, Ukraine, Kansas, New Jersey, Yugoslavia

***Penthimia picta* Provancher, 1872**

Penthimia picta Provancher, 1872: 352

Distribution: Bangladesh (new record), Germany, Japan, Russia, Albania, Austria, Azerbaijan, Belgium, Bohemia, Bulgaria, Czech Republic, Czechoslovakia, Russia, France, Germany, Great Britain, Greece, Hungary, Italy, Japan, Latvia, Macedonia, Moldova, Moravia, Netherlands, Poland, Portugal, Romania, Serbia, Slovakia, Spain, Switzerland, Syria, Tunisia, Turkey, Ukraine, Kansas, New Jersey, Yugoslavia.

Tribe Scaphoideini

Scaphoidella sp. Vilbaste, 1968

Scaphoidella sp. Vilbaste, 1968:133

Distribution: India, Thailand, China, Bangladesh (new record).

Tribe Idiocerini Baker, 1915

***Idioscopus clypealis* (Lethierry, 1889)**

Idiocerus clypealis Lethierry, 1889: 252

Distribution: India, Pakistan, Taiwan, southern China, Philippines, Malaysia, Indonesia, Brunei, Australian.

***Rhytidodus decimusquartus* (Schrank, 1776)**

Cicada decimusquartus Schrank, 1776: 76

Distribution: Argentina, Bangladesh (new record).

Infraorder Fulgoromorpha

Family Fulgoridae

Tribe Laternarini

***Pyrops spinolae* (Westwood, 1842)**

Synonym:

Fulgora spinolae Westwood, 1842: 118

Hotinus spinolae (Westwood, 1842)

Hotinus nigrirostris Walker, 1858

Fulgora nigrirostris (Walker, 1858)

Laternaria spinolae (Westwood, 1842)

Pyrops condorinus (Lallemand, 1960)

Laternaria nigrirostris (Walker, 1858)

Distribution: Bangladesh, India, Cambodia, Vietnam.

***Pyrops lathburii* (Kirby, 1818)**

Synonym:

Fulgora lathburii Kirby, 1818: 450

Flata lathburii (Kirby, 1818)

Hotinus lathburi (Kirby, 1818)

Laternaria lathburii (Kirby, 1818)

Distribution: Bangladesh, China, Vietnam

Tribe Limoisini

***Limois westwoodii* (Hope, 1843)**

Synonym:

Lystra westwoodi Hope, 1843 (sic): 133

Lystra westwoodii Hope, 1843

Limois westwoodi (Hope, 1843) (sic)

Distribution: Bangladesh

Tribe Aphaenini

***Penthicodes (Penthicodes) farinosa farinosa* (Weber, 1801)**

Synonym:

Lystra dimidiata Hope, 1843

Aphana dimidiata (Hope, 1843)

Aphaena scutellaris White, 1846

Aphaena farinosascutellarum White, 1846

Penthicodes scutellaris (White, 1846)

Aphaena farinosa var. *scutellaris* White, 1846

Penthicodes (Penthicodes) farinosa (Weber, 1801) : 21

Distribution: Bangladesh

***Penthicodes (Ereosoma) caja* (Walker, 1851)**

Synonym:

Aphaena caja Walker, 1851: 278

Aphana caja (Walker, 1851)

Penthicodes caja (Walker, 1851)

Distribution: Bangladesh, India, Myanmar, Laos, Thailand, Vietnam, Malaysia.

***Lycorma imperialis* (White, 1846)**

Synonym:

Aphana imperialis White, 1846: 330-333

Aphaena imperialis (White, 1846)

Lycorma imperialis (White, 1846)

Aphaena placabilis Walker, 1858

Lycorma imperialis placabilis (Walker, 1858)

Distribution: Bangladesh

***Lycorma imperialis punicea* (Hope, 1843)**

Synonym:

Lystra punicea Hope, 1843: 133

Aphana punicea (Hope, 1843)

Aphaena delectabilis Walker, 1858

Lycorma punicea (Hope, 1843)

Distribution: Bangladesh

***Lycorma delicatula* (White, 1845)**

Synonym:

Aphaena delicatula White, 1845: 37

Lycorma delicatulum (White, 1845)

Distribution: Bangladesh, China, South Korea, Japan, Vietnam

***Aphaena (Aphaena) aurantia* (Hope, 1840)**

Synonym:

Aphana aurantia Hope, 1840: 443

Aphaena aurantia (Hope, 1840)

Aphana aurora Hope, 1843

Aphaena aurora (Hope, 1843)

Euphria aurantia (Hope, 1840)

Euphria aurora (Hope, 1843)

Distribution: Bangladesh, India, China

***Aphaena (Callidepsa) amabilis* (Hope, 1843)**

Synonym:

Aphana amabilis Hope, 1843: 132

Aphaena amabilis (Hope, 1843)

Euphria amabilis (Hope, 1843)

Aphaenalectissima Walker, 1858

Euphrialectissima (Walker, 1858)

Distribution: Bangladesh, Vietnam

***Polydictya basalis* (Hope, 1843)**

Synonym:

Eurybrachis basalis Hope, 1843: 134

Polydictya basali (Hope, 1843)(sic)

Distribution: Bangladesh, Myanmar, Vietnam.

Tribe Eurybrachini

***Messena pulverosa* (Hope, 1843)**

Synonym:

Eurybrachys pulverosa Hope, 1843: 134

Distribution: Bangladesh

***Purusha reversa* (Hope, 1843)**

Synonym:

Eurybrachys reversa Hope, 1843: 134

Messena reversa Hope, 1843

Distribution: Bangladesh, India

Tribe Loxocephalini

***Loxocephala aeruginosa* (Hope, 1840)**

Synonym:

Lystra aeruginosa Hope, 1840: 443

Loxocephala aeruginosa (Hope, 1840)

Eurybrachys aeruginosa (Hope, 1840)

Distribution: Bangladesh, India, Vietnam

Family Dictyopharidae

Tribe Dictyopharini

***Dictyophara asperae* Shakila, 1984**

Dictyophara asperae Shakila, 1984: 87

Distribution: Bangladesh

***Dictyophara sauropsis* Walker, 1862**

Synonym:

Dictyophora sauropsis Walker, 1862: 306

Dictyophara sauropsis Walker, 1862

Distribution: Bangladesh

***Raivuna sinica* Walker, 1851**

Dictyophara sinica Walker, 1851: 321

Distribution: Korea, Pakistan and Bangladesh (new record).

Tribe Lophopini

***Corethrura fuscovaria* Hope, 1843**

Synonym:

Corethrura fuscovaria Hope, 1843: 321

Lophops fuscovarius (Hope, 1843)

Distribution: Bangladesh

***Elasmoscelis perforata* Walker, 1862**

Synonyms:

Elasmoscelis perforata Walker, 1862:309

Elasmoscelis platypoda Kirby, 1891

Distribution: Bangladesh, China, Thailand, Taiwan, Japan, Indonesia.

Family Achilidae

Tribe Plectoderini

***Deferunda stigmatica* Distant, 1912**

Deferunda stigmatica Distant, 1912: 186

Distribution: Bangladesh

***Deferunda qiana* Chen and He, 2010**

Deferunda qiana Chen and He, 2010: 64

Distribution: Korea, China and Bangladesh (New record).

Family Cixiidae

Tribe Bothriocerini

***Borysthenes suknanicus* Distant, 1911**

Borysthenes suknanicus Distant, 1911: 735-747

Distribution: Bangladesh, India

Tribe Pentastirini

***Oliarus tectonae* Shakila, 1984**

Oliarus tectonae Shakila, 1984: 295

Distribution: Bangladesh

Tribe Cixiini

***Bangoliarus truncatus* Shakila, 1984**

Bangoliarus truncatus Shakila, 1984: 314

Distribution: Bangladesh

Family Delphacidae

Tribe Delphacini

***Harmalia cordata* Yang, 1989**

Harmalia cordata Yang, 1989: 210

Distribution: Bangladesh, Indonesia, Philippines, Thailand, Taiwan

***Nilaparvatalugens* (Stål, 1854)**

Synonym:

Delphax lugens Stål, 1854: 246

Delphax sordescens Motschulsky, 1863

Kalpa aculeata Distant, 1906

Delphax parysatis Kirkaldy, 1907

Delphax ordovix Kirkaldy, 1907

Liburnia sordescens Melichar, 1903

Delphacodes anderita Muir, 1917

Nilaparvata sordescens Muir, 1922

Distribution: Bangladesh, Brunei, China, Cambodia, Japan, India, Myanmar, Indonesia, Korea, Malaysia, Nepal, Laos, Sri Lanka, Pakistan, Vietnam, Thailand, Singapore, Philippines.

***Perkinsiella rattlei* Muir, 1911**

Perkinsiella rattlei Muir, 1911: 6

Distribution: China, India and Bangladesh.

***Sardia rostrata* (Kirkaldy, 1906)**

Sardia rostrata (Kirkaldy, 1906): 96

Distribution: Australia, Bangladesh, China, France, Iran and Sri-Lanka.

***Tagosodes cubanus* (Crawford, 1914)**

Dicranotropis cubanus Crawford, 1914: 595

Distribution: Bangladesh (new record), USA, Belize, Bolivia, Brazil, Colombia, Costa Rica, Cuba, Jamaica, Martinique, Mexico, Nicaragua, Panama, Peru, Puerto Rico, Bermuda, Trinidad, Venezuela; Nigeria, Ivory Coast.

***Opiconsiva albicollis* (Motschulsky, 1863)**

Synonyms:

Delphax albicollis De Motschulsky, 1863: 110

Liburnia albicollis De Motschulsky, 1863

Kakuna sameshimai Matsumura & Ishihara, 1945

Harmalia sameshimai (Matsumura & Ishihara, 1945)

Toya yanoi (Ishihara, 1952)

Opiconsiva koreacola Kwon, 1982

Corbulokorea cola Kwon, 1982

Distribution: Bangladesh, Japan, Madagascar, Sri Lanka, South Korea, Philippines, Taiwan, Turkey, USA.

***Opiconsiva tangira* (Matsumura, 1910)**

Synonym:

Delphax tangira Matsumura, 1910: 30

Toya tangira (Matsumura, 1910)

Corbulo tangira (Matsumura, 1910)

Opiconsiva tangira (Matsumura, 1910)

Opiconsiva nigra (Ding & Tian, 1980)

Distribution: Bangladesh, India, Morocco, Thailand, Sri Lanka.

***Sogatella furcifera* (Horváth, 1899)**

Synonym:

Delphax furcifera Horváth, 1899: 372

Liburnia furcifera (Horváth, 1899)
Sogata pallescens Distant, 1912
Megamelus albolineosus Crawford, 1914
Opiconsiva gloriosa Distant, 1917
Opiconsiva insularis Distant, 1917
Opiconsiva balteata Distant, 1917
Opiconsiva colorata Distant, 1917
Opiconsiva derelicta Distant, 1917
Delphacodes albolineosa Osborn, 1926

Distribution: Bangladesh, Brazil, Australia, Cambodia, India, Thailand, Myanmar, Nepal, Bhutan, China, Egypt, Fiji, Florida, Great Britain, Indonesia, Iran, Israel, Italy, Jamaica, Japan, Kenya, Korea, Malaysia, Laos, Mexico, Mongolia, Nigeria, Pakistan, Philippines, Russia, Saudi Arabia, South Africa, Taiwan, USA, Spain, Vietnam.

Tribe Tropidocephalini

***Tropidocephala festiva* Distant, 1906**

Tropidocephala festiva Distant, 1906: 478

Distribution: Bangladesh (new record), China, Indonesia, Japan, Malaysia, Philippines, Sri Lanka, Taiwan.

Family Issidae

***Pinocchias natus* Gnezdilov & Wilson, 2005**

Pinocchias natus Gnezdilov & Wilson, 2005: 22

Distribution: Bangladesh

Family Ricaniidae

***Ricania amygdalusae* Shakila, 1984**

Ricania amygdalusae Shakila, 1984: 266

Distribution: Bangladesh

***Ricania zebra* Distant, 1906**

Ricania zebra Distant, 1906: 379

Distribution: Bangladesh

***Ricanoptera fenestrata* (Fabricius, 1803)**

Synonym:

Ricania mellerborgi Stål, 1854

Ricanoptera mellerborgi (Stål, 1854)

Flata fenestrata Fabricius, 1803: 51

Ricania fenestrata Schaum, 1850

Flatoides fenestrata Walker, 1851

Flatoides discalis Walker, 1857

Distribution: Bangladesh, Indonesia, Malaysia, Myanmar, Philippines, Singapore

Family Tettigometridae

***Egropa bengalensis* Distant, 1909**

Egropa bengalensis Distant, 1909: 41

Distribution: Bangladesh, India

Family Derbidae

Tribe Cedusini

***Hauptenia bandarbanensis* Jhan and Rahman, 2016**

Hauptenia bandarbanensis Jhan and Rahman, 2016: 2

Distribution: Bangladesh.

References

Bierman CJH (1910) Homopteren aus Niederländisch Ost-Indien. II herausgegeben von D. MacGillavry und K. W. Dammerman. *Notes from the Leyden Museum*, 33:1-68.

Bloem JH, Duffels JP (1976) The *Dundubia* terpsichore group of the genus *Dundubia* Amyot and Serville, 1843. *Bulletin Zoölogisch Museum, Universiteit van Amsterdam*, 5:141-154.

Chen XS, He TT (2010) Two new species of genus *Deferunda* Distant (Hemiptera: Fulgoromorpha: Achilidae) from southwest China. *Zootaxa*, 2335: 59-68.

Constant J (2010) The lanternfly genus *Penthicodes*: key to the species and review of the "Ereosoma group" with two new species and one new subspecies (Hemiptera: Fulgoromorpha: Fulgoridae). *Zootaxa*, 2523: 1-26.

Crawford DL (1914) A contribution toward a monograph of the Homopterous insects of the family Delphacidae of North and South America. *Proceedings of the United States National Museum*. Washington, 46: 557-640.

De Motschulsky VI (1859) "Homoptères." In *Insectes des Indes orientales, et de contrées analogues. Étud. Entomol.*, 8: 25-118.

De Motschulsky VI (1863) Essai d'un catalogue des insectes de l'île de Ceylan. *Bulletin de la Société Impériale des Naturalistes de Moscou*. Moscou, 36: 1-153.

Distant WL (1881) Descriptions of new species belonging to the Homopterous Family Cicadidae. *Transactions of the Royal Entomological Society of London*, 627-648.

Distant WL (1888a) Descriptions of new species of Oriental Homoptera belonging to the family Cicadidae. *Ann. Mag. Nat. Hist.* (6), 1: 291-298.

Distant WL (1888b) Viaggio di Leonardo Fea in Birmania e regioni vicine. VIII. Enumeration of the Cicadidae collected by Mr. L. Fea in Burma and Tenasserim. *Ann. Mus. Genova*. (2)I, 6: 453-459.

Distant WL (1908) Rhynchota. IV. Homoptera and appendix (Pt.). *The fauna of British India, including Ceylon and Burma*. London: Taylor & Francis. XV+501 p.

Distant WL (1906) The fauna of British India, including Ceylon and Burma. 3: 503 pp. Lt. Col. C. T. Birgham.

Distant WL (1909) Descriptions of two new species of Rhynchota from Bengal. *Annals and Magazine of Natural History*. London (Ser. 8), 3: 40-42.

Distant WL (1911) Descriptions of new genera and species of Oriental Homoptera. *Annals and Magazine of Natural History*. London (Ser. 8), 8: 735-747.

Distant WL (1912) Descriptions of new genera and species of Oriental Homoptera. *Annals and Magazine of Natural History*. London. (Ser. 8), 9: 181-194.

- Drury D (1773) Illustrations of natural history, wherein are exhibited upwards of two hundred and twenty figures of exotic insects, according to their different genera; very few of which have hitherto been figured by any author, ...many of them. 2: 1-90. pl. 1-50.
- Fabricius JC (1803) Rhyngota. In: Fabricius JC. 1803 *Systema Rhyngotorum secundum ordines, genera, species, adiectis synonymis, locis, observationibus, descriptionibus*, p. 1-101.
- Gnezdilov VM, Wilson MR (2005) New genera and species of the tribe Parahiraciini (Hemiptera: Fulgoroidea: Issidae). *Acta Entomologica Slovenica*, 13(1): 21-28.
- Goeze JAE (1778) Hemiptera. In *Entomologische beyträge zu des ritter Linné Zwolften ausgabe des natursystems. Bey Weidmanns erben und reich. Leipzig. Vol./pp.: 2:i-lxxii*, 1-352.
- Hope FW (1840) Descriptions of some new insects, collected in Assam by William Griffith, Esq., Assistant-Surgeon in the Madras Medical Service, and attached to the late Scientific Mission to Assam. *The Transactions of the Linnean Society of London. Second series. Zoology*, 18: 435-447.
- Hope FW (1843) On some rare and beautiful insects from Silhet, chiefly in the collection of Frederick John Parry, Esq. F. L. S. *The Transactions of the Linnean Society of London. London* 19: 131-136].
- Horváth G (1899) Hémiptères de l'île de Yesso (Japon). *Természetráji Füzetek. Budapest*, 22: 365-374.
- Jhan PK, Rahman MA, Khan MMH, Jahan SMH (2016) A newly recorded genus *Hauptenia* Szwedo (Hemiptera: Fulgoromorpha: Derbidae) in Bangladesh, with description of a new species. *International Journal of Innovative Research*, 1(1): 1-4.
- Kirby WF (1818) A century of insects, including several new genera described from his cabinet. *The Transactions of the Linnean Society of London. Second series. Zoology*, 12: 375-453.
- Kirkaldy GW (1909) Hemiptera, old and new, No. 2. *The Canadian Entomologist*, 41: 388-392.
- Lee YJ (2016) Erections of *Changa* gen. nov. and *Songga* gen. nov. of Aolina (Hemiptera: Cicadidae: Cicadinae: Dundubiini) with taxonomic notes and a synonymization. *Zootaxa*, 4084(4): 596-600.
- Lethierry LF (1889) Definitions of three new Homoptera. *Journal of the Asiatic Society of Bengal*, 58: 252-253.
- Linnaeus C (1758) *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus differentiis, synonymis, Locis. Tomus I. Holmiae. Laurentii Salvii*. 823 pp.
- Matsumura S (1914) Die Jassinen und einige neue Acocephalinen Japans. *The journal of the Sapporo Agricultural College, Sapporo, Japan.*, 5(7): 165-240.
- Matsumura S (1910) Neue Cicadinen aus Europa und Mittelmeergebiet (Schluss). *Journal of the College of Sciences, Imperial University of Tokyo. Tokyo*, 27(18): 1-38.
- McKamey SH (2006) Further new genus-group names in Cicadellidae (Hemiptera). *Proc. Entomol. Soc. Washington.*, 108(3): 502-510.
- Melichar L (1903) Homopteren-Fauna von Ceylon. 1903: 248 pp.
- Muir FAG (1911) On some new species of leafhopper (*Perkinsiella*) on sugar cane. Bulletin. Hawaiian Sugar Planters' Association Experiment Station. Division of Entomology, 9: 4-11.
- Provancher L (1872) Description de plusieurs Hémiptères nouveaux. *Le Naturaliste Canadien.*, 4:350-352.
- Rao KR (1989) Descriptions of some new leafhoppers (Homoptera: Cicadellidae) with notes on some synonymies and imperfectly known species from India. *Hexapoda*, 1: 59-83.
- Shakila M (1984) *Morphotaxonomical studies of some families of Fulgoroidea found in Pakistan and adjoining countries (Homoptera: Fulgoroidea)*. Thesis. 387 pp. University of Karachi, Department of Zoology, Karachi (Oregon).
- Stål C (1865) Homoptera nova vel minus cognita. Öfversigt af Kongliga Svenska Vetenskaps-Akademiens Förhandlingar., 22:145-165
- Stål C (1870) Hemiptera insularum Philippinarum. Bidrag till Philipponska öarnes Hemipterfauna. Öfversigt af Kongliga Svenska Vetenskaps-Akademiens Förhandlingar., 27:607-776.
- Stål C (1854) Nya Hemiptera. Öfversigt af Kongliga Svenska Vetenskaps-Akademiens Förhandlingar. *Stockholm*, 11: 231-255.
- Uhler PR (1896) Summary of the Hemiptera of Japan presented to the United States National Museum by Professor Mitzukuri. *Proceedings of the United States National Museum*, 19:255-297.
- Vilbaste J (1968) Systematic treatise of cicadas found on the edge of the coastal regions. Über die Zikadenfauna des Primorje Gebietes. Valgus, Tallin, 195 pp.
- Walker F (1850) List of the specimens of homopterous insects in the collection of the British Museum. (Published by) *Order of Trustees. London*, 1: 1-260.
- Walker F (1851) List of the specimens of homopterous insects in the collection of the British Museum. *British Museum, London*, Part III: (4): 637-907.
- Walker F (1850) List of the specimens of Homopterous insects in the collection of the British Museum. Part I. Edward Newman; London: 260.
- Walker F (1858) Homoptera. In *Insecta saundersiana: or characters of undescribed insects in the collection of William Wilson Saunders, Esq. John Van Voorst. London*. 117 p.

- Walker F (1851) List of the specimens of Homopterous insects in the collection of the British Museum. 2: 261-636.
- Walker F (1862) Characters of undescribed species of Homoptera in the collection of F. P. Pascoe, F. L. S. *The Journal of entomology: descriptive and geographical* 1: 303-319.
- Weber F (1801) Rhyngota. In: Weber F. 1801 – *Observationes entomologicae, continentes novorum quae condidit generum characteres, et nuper detectarum specierum descriptiones*, 1801. p. 112-115.
- Westwood JO (1842) Insectorum novorum centuria, auctore. *Annals and Magazine of Natural History* 9: 118-119.
- Westwood JO (1839) On the entomology of the Himalayas and of India. *Illustrations of the botany and other branches of the natural history of the Himalayan Mountains, and of the flora of Cashmere by J. Forbes Royle*, 1: i-lxxiv, 1-472.
- Westwood JO (1842) Insectorum novorum centuria, auctore. *Annals and Magazine of Natural History*. London 9: 118-119.
- White A (1845) Descriptions of a new genus and some new species of Homopterous Insects from the East in the collection of the British Museum. *Annals and Magazine of Natural History*. London 15: 34-37.
- White A (1846) Descriptions of some apparently new species of Homopterous insects in the collection of the British Museum. *Annals and Magazine of Natural History*. London, 17: 330-333.
- Yang CT (1989) Delphacidae of Taiwan II. *National Science Council (NSC), Special publication*, 6: 1-334.